MAPP No. 18850 PCS No. 04731

ROUNDUP[®] Ultra kills most garden weeds with a single application. This product is a home garden weedkiller for use on unwanted vegetation including trees and freshly cut tree stumps and areas not intended to bear vegetation (e.g. weeds growing through all hard surfaces including drives, patios, paths, tarmac, gravel and paving).

INSTRUCTIONS FOR USE

FOR USE ONLY AS AN AMATEUR WEEDKILLER. READ SAFETY INFORMATION INSIDE BEFORE USE.

SAFETYINSTRUCTIONS

Protection during/in use: If swallowed, seek medical advice immediately and show this container or label. WASH HANDS AND EXPOSED SKIN AFTER USE. KEEP OFF SKIN. WASH OFF SPLASHES IMMEDIATELY. Environmental protection: DO NOT CONTAMINATE WATER WITH PRODUCT OR ITS CONTAINER. Storage and disonsale KEEP PROINITIC AWAY FROM CHILD READ AND PETS KEEP AWAY FROM FOOD AND DRINK KEEP.

Storage and disposal KEEP PRODUCT AWAY FROM CHILDREN AND PETS. KEEP AWAY FROM FOOD AND DRINK. KEEP PRODUCT IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RISK & SAFETY INFORMATION

TO AVOID RISKS TO HUMAN HEALTH AND THE ENVIRONMENT, COMPLY WITH THE INSTRUCTIONS FOR USE.

• Read carefully and follow all instructions.

• Keep product away from food and drink.

If medical advice is needed, have product container or label at hand.

Keep product in original container.

 For UK only: Dispose of contents/container to a household waste recycling centre as hazardous waste except for empty containers which can be disposed of in the dustbin. Contact your local council for details.

For Rol only: Dispose of contents/container to a household waste recycling centre as hazardous waste except for triple

rinsed empty containers, which can be disposed of by recycling. Contact your local council for details.

Poisons information: For information or to report a poisoning incident in Ireland, contact the National Poisons Information Centre (01 809 2166), retain the label for reference



ROUNDUP' Ultra is a soluble concentrate containing 360 g/l (28.47 ‰/w) glyphosate acid (present as 441 g/l (34.86 ‰/w) potassium salt of glyphosate).



LAWN DESTRUCTION

HOW TO USE (continued)

Apply ROUNDUP[®] Ultra to actively growing lawn before the onset of dormancy, usually between May and September. Allow at least 7-10 days for ROUNDUP[®] Ultra to be translocated to the roots of treated lawn. Remove dying and deal awn. Either e-seed with grass seed or re-turd according to seed or turd providers instructions. ROUNDUP[®] Ultra should not be used on areas of lawn unless the user wants to kill them

FRESHLY CUT TREE AND SHRUB STUMPS

Cut the tree or shrub stem close to the ground in November to March, and thoroughly treat the cut surface of the stump as soon as possible after felling. Use a paint brush to apply the diluted ROUNDUP Ultra on to the cut stump. Ensure the whole cut surface is saturated, particularly around the edges and bark of the stump. Ensure the brush is thoroughly cleaned before and after use. Three stump dilutions: 20% solution in water (e.g. a mix of 20 ml of ROUNDUP⁹ Ultra and 80 ml water).

STANDING TREES

In November to March, use an axe to cut angled notches that completely penetrate the bark of standing trees. Cut at approximately knee height, one notch for trees up to 10 cm in diameter and use two or three notches in trees over 10 cm diameter. Using a X or 1 inch standard paint brush, apply 2 ml (thorough wetting of the inside of the notch) per 10 cm trunk diameter of neat ROUNDUP² UItra to each

cut. Do not treat in the period of active sap flow in the spring/early summer. Users should be aware that dead standing trees are a potential hazard and therefore should be felled before they fall over. IAPANESE KNOTWEED

Apply as a foliar spot application using a 2% solution (20 ml made up to 1 litre of water). For best results apply when the plant is flowering (usually August/September) but before the autumn dieback. Good coverage is essential. Spray the underside as well as the upper surface of the leaves. Repeat treatment may be necessary if regrowth appears. If Japanese Knotweed is growing amongst wanted plants apply 5-10 ml of a 20% (200ml ROUMDUP' Ultra made up to 1 litre of water) solution with water per stem

using the following method. After mid August but before leaf fall, cut the stems of the plants approximately 200mm above the base of the cane and 40 mm above a node. Rupture the central stem tissue with a screwdriver and apply 5-10 ml of a 20% solution of ROUNDUP 'Ultra into the hollow stem within 15 minutes of cutting using the 10 ml disposable pipette supplied with the pack. Gently dispense the einette avoiding solash back. Stems must be larger than 8mm diameter.

RESTRICTIONS

 If not applied as a spot treatment, the maximum yearly dose must not exceed the dose indicated in the above crop tables.

 Do not walk or allow children and pets into treated areas until the product has dried on leaves to avoid transfer to lawns and other desired vegetation.

 Maximum individual dose: 1 per year. When applied via pipette the maximum concentration must not exceed 200 ml of product per 1 litre of water. When applied as a spray the maximum concentration must not exceed 20 ml of product per 1 litre of water. When used on areas not intended to bear vegetation the maximum dose must not exceed 32 ml/30 m². When applied by injection for chemical thinning of trees the maximum concentration must not exceed 2 ml product/10 cm of tree diameter. When applied to tree stumps the maximum concentration must not exceed 200 ml of product per 1 litre of water.

Replanting and sowing

Light cover of annual weeds:	Replanting/sowing can commence the day after treatment
Dense cover, including perennial weeds:	Leave at least 7 days for the weedkiller to move to underground roots and shoots before digging or cultivating soil

perennial weeds: roots and shoots before digging or cultivating soil After application, rinse the sprayer three times, pouring rinse water over bare ground. Triple rinse empty bottle and pour rinse water over bare ground.				

FIRST AID

Eye contact: Immediately flush with plenty of water. If easy to do, remove contact lenses. Skin contact: Take off contaminated clothing, wristwatch, jewellery. Wash clothes and clean shoes

before re-use. Inhalation: Remove to fresh air.

Ingestion: Immediately offer water to drink. Never give anything by mouth to an unconcious person. DO NOT induce vomiting unless directed by medical personnel. If symptoms occur, get medical attention.

ROUNDUP[®] Ultra

ROUNDUP' Ultra is a non-selective, translocated weedkiller that is absorbed by leaves and freshly cut, woody surfaces and moves throughout the entire plant, above and below ground, to control annual and perennials weeds, including grasses and deep rooted perennials weeds. It also controls standing trees, newly cut tree stumps, tree spatings, brannelse, bracken, nettles, Japanese knotweed and other tough and woody weeds. ROUNDUP' Ultra is degraded by micro-organisms in the soil.

INSTRUCTIONS FOR USE FOR USE ONLY AS AN AMATEUR WEEDKILLER.

Apply as a foliar spray to weeds and saplings at any time when the weeds are actively growing and have adequate leaf area to absorb the spray. Freshly cut tree stumps and standing trees should be treated from November to March, to avoid the period of active sap flow in spring/early summer. Avoid spraying when rain is imminent. Rain within 6 hours of spraying may reduce effectiveness, so repeat treatment may be necessary. Avoid spraying in windy conditions, as spray drift to lawns and other wanted plants will damage or destroy them. Avoid all other spray contact with cultivated plants. If cultivated plants are treated by missake. IMMEDIATELY AND THOROUGHIY wash off the weekfiller with annele water.

WHERE TO USE

 ROUNDUP¹ Ultra controls annual weeds and most perennial weeds and trees around the garden and on areas not intended to bear vegetation such as hard surfaces, patios, drives, tarmac, paths and gravel areas it can be used to clean up weedy ground before planting vegetables, flowers, bubses and trees, and to renovate overgrown areas. Direct spray away from ponds and other surface water bodies.
 DO NOT USE ON WANTED PLANTS. DO NOT USE ON AREAS OF LAWN UNLESS YOU WANT TO KILL IT. DO NOT USE UNDER POLYTHENE OR GLASS.

HOW TO USE

 Although not a requirement, as good practice, we recommend wearing waterproof gloves and gardening footwar. Apply by sprayer to lightly wet the leaves. If thre is approximately 2 pints. S fittres is approximately I gallon. When diluted in 1 litre of water as surface of about 30 m⁻ can be treated. Do not apply through, or store in, galvanised or unlined steel sprayers. • Weeds show effects within 1-4 weeks. Dead or dying weeds can be dug in or composted without harming future plants.
 Annilai ADN PERINNIAI WEFRS [foliar anolications]

ROUNDUP [®] Ultra to be added to: Sprayer - 1 litre of water		Area treated	When to treat			
Annual weeds: Chickweed, Speedwell, Annual Meadow Grass	10 ml	30 m²	When actively growing, usually February – November			
Perennial grasses: including Couch	15 ml	30 m²	When actively growing, usually February – November			
Tough perennial deep rooted and broad-leaved weeds: Bindweed, Dandelions, Docks, Ground Elder*, Thistles*	15 ml	30 m²	Dandelions and Docks: when actively growing, usually April - Octol Bindweed, Ground Elder, Thistles: around the time of flowering			
Areas not intended to bear vegetation	15 ml	30 m²	When actively growing, usually February – November			
Lawn destruction	20 ml	30 m²	Treat in a period of active growth			
BRUSH WEEDS AND SAPLINGS (foliar applications)						
ROUNDUP [®] Ultra to be added to:		Area	Best time of year to treat			
Sprayer - 1 litre of water		treated				
	10 ml	treated 30 m²	Brambles, Nettles, Scrub: June - Septeml Horsetail, Braken: July - August Fully grown Nettles: cut back in early summer and treat regrowth in Septeml			
Sprayer - 1 litre of water Brush weeds: Brambles, Nettles,	10 ml 30 ml		Horsetail, Braken: July – August Fully grown Nettles: cut back in early			